

4. Heath CA, Cooper SA, Murray K, Lowman A, Henry C, MacLeod MA, et al. Validation of diagnostic criteria for variant Creutzfeldt-Jakob disease. *Ann Neurol*. 2010;67:761–70.
5. Kaski D, Mead S, Hyare H, Cooper S, Jampana R, Overell J, et al. Variant CJD in an individual heterozygous for PRNP codon 129. *Lancet*. 2009;374:2128. [http://dx.doi.org/10.1016/S0140-6736\(09\)61568-3](http://dx.doi.org/10.1016/S0140-6736(09)61568-3)
6. Krasemann S, Neumann M, Geissen M, Bodemer W, Kaup FJ, Schulz-Schaeffer W, et al. Preclinical deposition of pathological prion protein in muscle of experimentally infected primates. *PLoS ONE*. 2010;5:e13906. <http://dx.doi.org/10.1371/journal.pone.0013906>
7. Glatzel M, Abela E, Maissen M, Aguzzi A. Extraneural pathologic prion protein in sporadic Creutzfeldt-Jakob disease. *N Engl J Med*. 2003;349:1812–20. <http://dx.doi.org/10.1056/NEJMoa030351>
8. Peden AH, Ritchie DL, Head MW, Ironside JW. Detection and localization of PrP^{Sc} in the skeletal muscle of patients with variant, iatrogenic, and sporadic forms of Creutzfeldt-Jakob disease. *Am J Pathol*. 2006;168:927–35. <http://dx.doi.org/10.2353/ajpath.2006.050788>
9. Asante EA, Linehan JM, Gowland I, Joiner S, Fox K, Cooper S, et al. Dissociation of pathological and molecular phenotype of variant Creutzfeldt-Jakob disease in transgenic human prion protein 129 heterozygous mice. *Proc Natl Acad Sci U S A*. 2006;103:10759–64. <http://dx.doi.org/10.1073/pnas.0604292103>
10. Glatzel M, Pekarik V, Luhrs T, Dittami J, Aguzzi A. Analysis of the prion protein in primates reveals a new polymorphism in codon 226 (Y226F). *Biol Chem*. 2002;383:1021–5. <http://dx.doi.org/10.1515/BC.2002.109>
11. Trifilo MJ, Yajima T, Gu Y, Dalton N, Peterson KL, Race RE, et al. Prion-induced amyloid heart disease with high blood infectivity in transgenic mice. *Science*. 2006;313:94–7. <http://dx.doi.org/10.1126/science.1128635>
12. Ashwath ML, Dearmond SJ, Culclasure T. Prion-associated dilated cardiomyopathy. *Arch Intern Med*. 2005;165:338–40. <http://dx.doi.org/10.1001/archinte.165.3.338>
13. Herzog C, Sales N, Etcheagaray N, Charbonnier A, Freire S, Dormont D, et al. Tissue distribution of bovine spongiform encephalopathy agent in primates after intravenous or oral infection. *Lancet*. 2004;363:422–8. [http://dx.doi.org/10.1016/S0140-6736\(04\)15487-1](http://dx.doi.org/10.1016/S0140-6736(04)15487-1)
14. Tennent GA, Head MW, Bishop M, Hawkins PN, Will RG, Knight R, et al. Disease-associated prion protein is not detectable in human systemic amyloid deposits. *J Pathol*. 2007;213:376–83. <http://dx.doi.org/10.1002/path.2240>
15. Herzog C, Riviere J, Lescoutra-Etcheagaray N, Charbonnier A, Leblanc V, Sales N, et al. PrPTSE distribution in a primate model of variant, sporadic, and iatrogenic Creutzfeldt-Jakob disease. *J Virol*. 2005;79:14339–45. <http://dx.doi.org/10.1128/JVI.79.22.14339-14345.2005>

Address for correspondence: Markus Glatzel, Institute of Neuropathology, University Medical Center Hamburg-Eppendorf, Martinistrasse 52, D-20246 Hamburg, Germany; email: m.glatzel@uke.de

etymologia

Shewanella haliotis

[shoo"ə-nel'ə hā"lī-o'tīs]

From the Greek halios (marine) and ōtos (ear), abalones, genus *Haliotis*, were first mentioned ≈2,500 years ago by Aristotle, who wrote of “the wild limpet (called by some the ‘sea ear’).” In D’Arcy Thompson’s translation of Aristotle, he notes that “wild limpet” is “commonly attributed to *Fissurella graecia* ... and conceals a forgotten name for *Haliotis*.” The “sea ear” was familiar to the Greeks and was named otia (little ear) by Pliny.

Shewanella haliotis, a species of rod-shaped, gram-negative, facultatively anaerobic bacteria, was first isolated from the gut microflora of abalones collected from the ocean near Yeosu, South Korea, by Kim et al. in 2007. The genus *Shewanella* had been previously named in 1985 by MacDonell and Colwell in honor of Scottish microbiologist James M. Shewan, for his work in fisheries microbiology.

Sources

1. Cox CW. California abalones, family Haliotidae. Sacramento (CA): The Resources Agency of California, Department of Fish and Game. Fish Bulletin no. 118; 1962.
2. Hunter R. The encyclopaedic dictionary: a new and original work of reference to all the words in the English language. Vol. IV, part 1. London: Cassell & Company, Limited; 1884.
3. Kim D, Baik KS, Kim MS, Jung BM, Shin TS, Chung GH, et al. *Shewanella haliotis* sp. nov., isolated from the gut microflora of abalone, *Haliotis discus hannai*. *Int J Syst Evol Microbiol*. 2007;57:2926–31. PubMed <http://dx.doi.org/10.1099/ijs.0.65257-0>
4. MacDonell MT, Colwell RR. Phylogeny of the Vibrionaceae, and recommendation for two new genera, *Listonella* and *Shewanella*. *Systematic and Applied Microbiology*. 1985;6:171–82. [http://dx.doi.org/10.1016/S0723-2020\(85\)80051-5](http://dx.doi.org/10.1016/S0723-2020(85)80051-5)

Address for correspondence: Ronnie Henry, Centers for Disease Control and Prevention, 1600 Clifton Rd NE, Mailstop E03, Atlanta, GA 30333, USA; email: boq3@cdc.gov

DOI: <http://dx.doi.org/10.3201/eid1906.ET1906>